



Contents lists available at [ScienceDirect](#)

## Neuroscience Letters

journal homepage: [www.elsevier.com/locate/neulet](http://www.elsevier.com/locate/neulet)



### Corrigendum

## Corrigendum to “A utricular origin of frequency tuning to low-frequency vibration in the human vestibular system?” [Neurosci. Lett. 451 (2009) 175–180]

Neil P.M. Todd<sup>a,\*</sup>, Sally M. Rosengren<sup>b</sup>, James G. Colebatch<sup>b</sup>

<sup>a</sup> Faculty of Life Science, Jacksons Mill, University of Manchester, Manchester M60 1QD, UK

<sup>b</sup> Prince of Wales Clinical School and Medical Research Institute, University of New South Wales, Randwick, Sydney, NSW 2031, Australia

In Fig. 1B the grey tracing for the VEMP (earlier p response) is for the medially accelerated or ‘contralateral’ side while for the OVEMP the lighter trace (n11) refers to the side laterally accelerated or ‘ipsilateral’ side.

For the OVEMP section of Fig. 2, ‘IPSI’ and ‘CON’ should be reversed.

DOI of original article: [10.1016/j.neulet.2008.12.055](https://doi.org/10.1016/j.neulet.2008.12.055).

\* Corresponding author at: Faculty of Life Science, Jacksons Mill, University of Manchester, Oxford Road, Manchester M60 1QD, UK. Tel.: +44 161 306 5770; fax: +44 161 306 3887.

E-mail address: [neil.todd@manchester.ac.uk](mailto:neil.todd@manchester.ac.uk) (N.P.M. Todd).